## Office of Medical Superintendent Rajindra Hospital, Patiala.

No.RHP/24/ 1595

**Notice Inviting Quotations** 

Date: 20/01/24

Sealed quotations are invited from the Local reputed firm/original manufactures/authorized dealers of Medical/General Supplier on or before dated 29/01/2024 up to 11.30 AM for the following items. Quotations will be opened on the same day as on 12.30 PM. In case last date is declared as holiday by the Govt the quotations will be opened on the next working day.

Sr.No	Items Names & Pack Size	Specification
	2. The runner of I her blice	·
1	T3 (96 Wells)	- Assay based upon ELISA method with antibody coated technology
		- Kit should be able to run on ELISA reader
		- Kit should be able to rull oil Elisa reader
		- The shelf life of the kit should be minimum of one year the reconstituted reagent after
		- Opening the kit, having maximum stability should be given consideration.
		- The kit with maximum linearity, high specificity and high sensitivity should be given preference.
		- The size of the kit / number of tests per kit is to be indicated.
		- Accurate Procedure(less tedious procedure) of the test should be provided & normal range should be provided
		- Kit should include - microlite plate with coated wells Strip holder
		Adhesive plate sealers     Washings should be minimum(less than 5)
		- Incubation period less than 2 hours
2	T4 (96 Wells)	-do-
3	TSH (96 Wells)	-do-
4	Vitamin D 3 (96 Wells)	-do-
5	LH (96 Wells)	-do-
6	FSH (96 Wells)	-do-
7	PROLACTIN (96 Wells) TESTOSTERONE (96	-do-
8	Wells)	-d0-
9	PSA (96 Wells)	-do-
10	CA-125 (96 Wells)	-do-
11	CEA (96 Wells)	-do-
12	CA-15.3 (96 Wells)	-do-
13	Alpha fetoprotein (AFP) (96 wells)	-do-
14	CA-19.9 (96 Wells)	-do-
15	Ferritin (96 Wells)	-do-
16	Insulin (96 Wells	-do-
17	Beta HCG (96 Wells)	-do-
18	Amylase (40x1ml)	Assay based upon kinetic method utilizing CI-PNP.
19	CPK-MB (20x1.1 ml)	Immunoinhibiton based kinetic method
		It should be a single step procedure
		Kit should be able to run on semi autoanalyser
		Kit having the maximum linearity high sensitivity and specificity should be given preference
		Maximum number of days of stability of the reconstituted reagent should be the main consideration
		Procedure of the test should be provided
20	LDH (20X1.1 ml)	Assay based upon kinetic DGKC method utilizing NADH.
		The reaction volume should be 500 micro liter
		Kit should be able to run on semi-autoanalyser.
		Maximum linearity and maximum number of days of stability of reconstituted reagents should be the consideration.
	1130	Procedure of the test should be provided.

	CPK TOTAL(CK-NAC) (15X1.1 ml)	Immunoinhibiton based kinetic method
1		It should be a single step procedure
		Kit should be able to run on semi autoanalyser
		Kit having the maximum linearity high sensitivity and specificity should be given preference
		Maximum number of days of stability of the reconstituted reagent should be the main consideration
		Procedure of the test should be provided
22	Phosphorous (2X50 ml)	Ammonium molybdate method Complex based technology
		It should be an end point method
		It should be a single step test The reaction volume should be 500 ul
		Kit should be able to run on semi-autoanalyser
		Kit showing the maximum linearity high sensitivity and high specificity
		should be given preference
		Maximum number of days of stability of re-constituted reagent should be
		the main consideration  Procedure of the test should be provided
23	Calcium (2X50 ml)	OCPC
	,	Complex based technology
		It should be an end point method
		It should be a single step test
		The reaction volume should be 500 ul  Kit should be able to run on semi-autoanalyser
		Kit showing the maximum linearity high sensitivity and high specificity
		should be given preference
		Maximum number of days of stability of re-constituted reagent should be
		the main consideration
		Procedure of the test should be provided
24	Micro Protein KITS for	Enzymatic method,
	CSF (1x50 ML)	It should be a single step test
		It should be an end point method  Kit should be able to run on semi-autoanalyser
		The reaction volume should be 500ul
		Maximum number of days of stability of reconstituted reagent should be the
		consideration
		Test kit having the maximum linearity should be given preference The size of the kit / number of tests per kit is to be indicated
		Procedure of the test should be provided.
25	Blood Urea KIT (2X1	Enzymatic METHOD (Berthelot)
	LITRE)	Autozyme Urea is a two reagent system using two step Procedure
		It should be an end point method  Kit should be able to run on Colorimeter / semi-autoanalyser.
		Maximum number of days of stability of reconstituted reagent should be the
		consideration
		Test kit having the maximum linearity should be given preference
		The size of the kit / number of tests per kit is to be indicated Procedure of the test should be provided
		Procedure of the test should be provided
26	Uric Acid (4X50 ML)	Enzymatic method, It should be an end point method
		Kit should be able to run on semi-autoanalyser and autoanalyser.
		Maximum number of days of stability of reconstituted reagent should be the
		consideration
		Test kit having the maximum linearity should be given preference
		The size of the kit / number of tests per kit is to be indicated Procedure of the test should be provided
27	Cholestrol 5x20ml	Phosphotugstic acid and magnesium chloride as precipitant
21	Cholestion 3x20iiii	Shelf life should be minimum of one year
		Procedure of the test should be provided.

		GPO-PAP based enzymatic method	
	Triglycerides 5x20ml	It should be an end point procedure	
		The reaction volume should be 500 ul	
		Kit should be able to run on semi autoanalyser and auto analyzer.	
		It should be a single step method	
		Kit having the maximum linearity high sensitivity and maximum number of days of stability of the reconstituted reagents should be	
		given preference	
20	LIDL Chalaster Section 1	Procedure of the test should be provided	
29	HDL-Cholestrol 5x20ml	Phospho tungistic acid method	
		It should be an end point procedure	
		The reaction volume should be 500 ul	
		Kit should be able to run on semi autoanalyser and auto analyzer.	
		It should be a single step method	
		Kit having the maximum linearity high sensitivity and maximum number of days of stability of the reconstituted reagents should be given preference	
		Procedure of the test should be provided	
30	SGOT 5x20	Assay based upon kinetic method recommended IFCC utilizing LDH-NADH coupled assay	
		Kit should be able to run on semi-autoanalyser and autoanalyser.  The reaction volume should be 500ul	
		Maximum number of days of stability of reconstituted reagent should be the	
		consideration	
		vi.Test kit having the maximum linearity should be given preference  The size of the kit / number of tests per kit is to be indicatedProcedure of the	
31	SGPT 5x20	test should be provided  Assay based upon kinetic method recommended IFCC utilizing LDH-NADH	
		coupled assay	
		Kit should be able to run on semi-autoanalyser and autoanalyser.  The reaction volume should be 500ul	
		Maximum number of days of stability of reconstituted reagent should be the	
		consideration	
		Test kit having the maximum linearity should be given preference The size of the kit / number of tests per kit is to be indicated	
		- Procedure of the test should be provided	
32	Glucose 2x200 ml	Enzymatic method	
		It should be an end point method	
		Kit should be able to run on semi-autoanalyser and autoanalyser.  Maximum number of days of stability of reconstituted reagent should be the	
		consideration	
		Test kit having the maximum linearity should be given preference The size of the kit / number of tests per kit is to be indicated	
		Procedure of the test should be provided	
33	Total Bilirubin 4x50 ml	Diazo based methodology	
		It should be a single step test It should be an end point method	
		Kit should be able to run on semi-autoanalyser	
		The reaction volume should be 500ul	
		Maximum number of days of stability of reconstituted reagent should be the consideration	
		Test kit having the maximum linearity should be given preference	
		The size of the kit / number of tests per kit is to be indicated	
	700	Procedure of the test should	
34	Ammonium Sulphate 500gm Disodium Phenyl Phosphate	1/(AK)	
35 36	Phenyl Dihydrogen Phospha	te (AR) 500gm per pack	
37	Trichloracetic acid (TCA) A	Trichloracetic acid (TCA) AR 100gm	
	A CONTRACTOR OF THE CONTRACTOR		

38	Trisodium Citrate (AR) 500gm per apck
39	Sodium Dihydrogen Phosphate (NAH2PO4) (AR)
40	Sulphur Powder AR 500gm per apck
41	Liquor Ammonia AR 500 ml per apck
42	Barium Chloride AR 500gm per apck
43	Fouchest's Regeants AR 500gm per apck
44	Ehrlich's Regeant AR 500gm per apck
45	Zinc Sulphate AR 500gm per apck
46	Starch soluble extra pure AR 500gm per apck
47	D- Fructose AR 500gm per apck
48	Glactose EXtra Pure AR 100gm per apck
49	Benedict's regeant AR 5 lrt/pack
50	DextroseAR 500gm per apck
51	Auto Pipettee (Variable Sizes) 10-100 & 100-1000 MICROLITRE Good Quality
52	Multi channnel (8-Channel) Auto Pipettee (Variable Sizes) For ELISA Test Good Quality
53	Plastic Test Tubes (13x75 mm) 500 Pieces per pack. Good Quality
54	Aliquots (1.5 ML) (500 Pieces per pack) Good Quality
55	Disposable gloves(loose) ( Size 6.5 &7.0 No. (50 pair per pack) Good Quality
56	disposable micro tips (10-100 microlitre) 1000 piece /pack Good Quality
57	Disposable micro (tips 200-1000 microlitre) 500pcs/ pack Good Quality
58	Sodium Hypochlorite Solution 10% 5 LTR Good Quality
59	Tissue Roll 6 pieces per pack G''ood Quality
60	Filter paper 500 pieces per pack Good Quality

ੀ ਕੁਟੇਸ਼ਨਾਂ ਕੰਪਿਊਟਰ ਰਾਹੀ ਟਾਈਪ ਹੋਣੀ ਚਾਹੀਦੀਆ ਹਨ। ਹੱਥ ਲਿਖਤੀ ਕੁਟੇਸ਼ਨਾਂ ਮੰਨਣਯੋਗ ਨਹੀ ਹੋਣਗੀਆ।ਕੁਟੇਸ਼ਨ ਵਿੱਚ ਰੇਟ ਬਿਨ੍ਹਾਂ ਜੀ.ਐਸ.ਟੀ. ਤੋ ਦੱਸਿਆ ਜਾਵੇਗਾ।

ਮੈਡੀਕਲ ਸੁਪਰਡੰਟ, ਰਾਜਿੰਦਰਾ ਹਸਪਤਾਲ ਪਟਿਆਲਾ ਪਾਸ ਕਿਸੇ ਵੀ ਕੁਟੇਸਨ/ਕੁਟੇਸ਼ਨਾ ਨੂੰ ਬਿਨ੍ਹਾਂ ਕੋਈ ਕਾਰਨ ਦੱਸੇ ਰੱਦ

ਕਰਨ ਦਾ ਪੂਰਾ–ਪੂਰਾ ਹੱਕ ਰਾਖਵਾਂ ਹੋਵੇਗਾ

2

3 ਕਿਸੇ ਵੀ ਕਿਸਮ ਦਾ ਝਗੜੇ ਸਬੰਧੀ Arbitration Act ਦੇ ਅਨੂਸਾਰ Sole Arbitrator ਡਾਇਰੈਕਟਰ ਪ੍ਰਿੰਸੀਪਲ, ਮੈਡੀਕਲ ਕਾਲਜ, ਪਟਿਆਲਾ ਹੋਣਗੇ ਅਤੇ ਉਹਨਾਂ ਦਾ ਫੈਸਲਾ ਮੱਨਯੌਗ ਅਤੇ ਅੰਤਿਮ ਹੋਵੇਗਾ।

ਰ ਫਰਮ ਕੋਲ ਜੀ.ਐਸ.ਟੀ, ਪੈਨ ਨੰਬਰ ਅਤੇ ਡੀ.ਐਲ. ਨੰਬਰ ( Wherever applicable) ਹੋਣਾ ਲਾਜ਼ਮੀ ਹੈ।

5 ਟੀ.ਡੀ.ਐਸ. ਜੀ.ਐਸ.ਟੀ. ਐਕਟ ਅਤੇ ਇੰਨਕਮ ਟੈਕਸ ਐਕਟ ਦੇ ਰੂਲਾ ਮੁਤਾਬਿਕ ਕੁੱਟਿਆ ਜਾਵੇਗਾ।

6 ਇਹਨਾ ਕੁਟੇਸ਼ਨਾ ਦੀ ਮਿਆਦ ਛੇ ਮਹੀਨੀਆ ਤੱਕ ਦੀ ਹੋਵੇਗੀ।ਜੇਕਰ ਸਰਕਾਰ ਵੱਲੋਂ ਕਿਸੇਪ੍ਰਕਾਰ ਦੀ ਪਾਲਿਸੀ ਨਿਰਧਾਰਿਤ ਕੀਤੀ ਜਾਂਦੀ ਹੈ ਤਾਂ ਉਹ ਸਬੰਧਤ ਫਰਮ ਤੇ ਲਾਗੂ ਹੋਵੇਗੀ।

7 ਉਕਤ ਮੰਗੀ ਗਈ ਕੁਟੇਸ਼ਨਾ ਇਸ ਦਫਤਰ ਦੇ ਰਸ਼ੀਟ ਸਾਖਾ ਵਿੱਚ ਰੱਖੇ ਗਏ ਸੀਲ ਬੰਦ ਡੰਬੇ ਵਿੱਚ ਸਮੇਂ ਸਿਰ ਪਾਇਆ ਜਾਣ।

ਮੈਡੀ**ਕੀ ਸੁੱ**ਧਰਡੰਟ ਰਾਜਿੰਦਰਾ ਹਸਪਤਾਲ ਪਟਿਆਲ੍ਹਾ।